

**Editorial Office**

Dr. Hiroyuki Tanaka (Editor-in-Chief)

Faculty of Agriculture, Tottori University, JAPAN

Dr. Shotaro Takenaka (Editor, Vice)

Department of Plant Life Science, Faculty of Agriculture, Ryukoku University, JAPAN

Dr. Goro Ishikawa (Editor, Associate)

NARO Institute of Crop Science, JAPAN

Dr. Kanako Kawaura (Editor, Associate)

Kihara Institute for Biological Research, Yokohama City University, JAPAN

Dr. Yoshihiro Matsuoka (Editor, Associate)

Department of Bioresources, Fukui Prefectural University, JAPAN

Dr. Hidetaka Nishida (Editor, Associate)

Graduate School of Environmental and Life Science, Okayama University, JAPAN

Dr. Tsuneo Sasanuma (Editor, Associate)

Department of Bioresource Engineering, Faculty of Agriculture, Yamagata University, JAPAN

Dr. Shigeo Takumi (Editor, Associate)

Laboratory of Plant Genetics, Graduate School of Agricultural Science, Kobe University, JAPAN

**Advisory Board**

Dr. Justin D. Faris

USDA-ARS Cereal Crops Research Unit, Northern Crop Science Laboratory, North Dakota, USA

Dr. Andreas Houben

Institute of Plant Genetics and Crop Plant Research (IPK), GERMANY

Dr. Shahryar F. Kianian

Cereal Disease Laboratory, USDA-ARS, Minnesota, USA

Dr. Masahiro Kishii

Global Wheat Program, International Maize and Wheat Improvement Center (CIMMYT), MEXICO

Dr. Evans Lagudah

CSIRO Plant Industry, AUSTRALIA

Dr. Hakan Ozkan

Department of Field Crops, Faculty of Agriculture, University of Cukurova, TURKEY

Dr. Hisashi Tsujimoto

Arid Land Research Center, Tottori University, JAPAN

Dr. Xueyong Zhang

Institute of Crop Sciences, Chinese Academy of Agricultural Sciences, Beijing, CHINA